



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07D 211/60, C07B 55/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/09290</b> <b>(43) International Publication Date:</b> 28 March 1996 (28.03.96)
<b>(21) International Application Number:</b> PCT/GB95/02247 <b>(22) International Filing Date:</b> 22 September 1995 (22.09.95) <b>(30) Priority Data:</b> 94306962.5                      23 September 1994 (23.09.94)      EP (34) <i>Countries for which the regional or</i> <i>international application was filed:</i> GB et al. 9504927.6                      10 March 1995 (10.03.95)      GB <b>(71) Applicant (for all designated States except US):</b> CHIRO- SCIENCE LIMITED [GB/GB]; Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DYER, Ulrich, Conrad [GB/GB]; Chiroscience Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). LOCK, Christo- pher, James [GB/GB]; Chiroscience Limited, Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). WOODS, Martin [GB/GB]; Chiroscience Limited, Cam- bridge Science Park, Milton Road, Cambridge CB4 4WE (GB). <b>(74) Agent:</b> GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).		<b>(81) Designated States:</b> AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> RACEMISATION AND ASYMMETRIC TRANSFORMATION PROCESSES USED IN THE MANUFACTURE OF LEVOBUPIVACAINE AND ANALOGUES THEREOF		
<b>(57) Abstract</b> <p>A process for the racemisation of an optically-enriched piperidine-2-carboxanilide compound, comprises heating the compound in the presence of an alkanolic or arylalkanoic acid. A process for the asymmetric transformation of such a compound comprises heating the compound in the presence of an acid as defined above, a chiral acid resolving agent and an inert cosolvent.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

RACEMISATION AND ASYMMETRIC TRANSFORMATION  
PROCESSES USED IN THE MANUFACTURE OF LEVOBUPIVACAINE  
AND ANALOGUES THEREOF

Field of the Invention

5           This invention relates to the racemisation and dynamic resolution of optically-enriched heterocyclic carboxanilides, e.g. piperidine-2-carboxanilides such as levobupivacaine.

Background of the Invention

10           Compounds of formula 1 (see formulae, below) wherein  $R^1$  is methyl, n-propyl, n-butyl or cyclopropyl and  $R^2$  is a 2,6-dimethylphenyl have utility as local anaesthetics. Biological studies have shown that (S)-enantiomers of such compounds display lower cardiotoxicity than the  
15           corresponding racemates whilst maintaining the same anaesthetic potency, and are therefore potentially more beneficial for clinical uses. Thus there is a requirement for efficient processes to manufacture compounds of formula 1 in the form of single enantiomers. For this purpose,  
20           conventional resolution approaches invariably afford up to 50% of the unwanted enantiomer. To improve atom utilisation in such processes, it is desirable to recycle the unwanted enantiomer by effecting its racemisation to provide material suitable for subsequent resolution.  
25           Additional benefits may be attainable by "asymmetric transformation", comprising simultaneous racemisation and crystallisation-induced resolution in a one-pot process.

          Fyhr et al, Acta Pharm. Suecica 25(3):121-132 (1988), disclose the racemisation of ropivacaine hydrochloride  
30           (1.HCl,  $R^1$  = n-propyl,  $R^2$  = 2,6-dimethylphenyl, absolute configuration = S) in dilute aqueous solution at pH 1-6, using HCl, and 80-130°C. The results are presented as a preformulation stability study and merely serve to indicate that ropivacaine racemises slowly in aqueous media.

35           Shiraiwa et al, Bull. Chem. Soc. Jpn. 64:3251-3255 (1991), disclose asymmetric transformation of 2-piperidine-carboxylic acid, by heating in an alkanolic acid solvent in

the presence of an chiral acid resolving agent and an aldehyde. The latter component is believed to assist racemisation by formation of a cationic Schiff base intermediate, a mechanistic pathway which can also operate  
5 on piperidine-2-carboxanilides 1 only in cases where  $R^1 = H$ . Again, this process is unsuitable for operation on a manufacturing scale, not least because it uses environmentally-unacceptable reagents.

#### Summary of the Invention

10 The present invention is based on the suprising discovery that N-heterocyclic-2-carboxanilides, including compounds of formula 1 wherein  $R^1$  is H, methyl, *n*-propyl, *n*-butyl or cyclopropyl and  $R^2$  is phenyl optionally substituted with one or more methyl groups, undergo rapid  
15 racemisation when heated in solution in the presence of a carboxylic acid  $R^3CO_2H$  wherein  $R^3$  is either *n*-alkyl or aryl (exemplified in Scheme 1) or any acid having a pKa of -1 to +6, relative to water. The reaction can be carried out in a wholly or substantially non-aqueous system, e.g. either  
20 in a solution of neat acid or in the presence of an inert cosolvent such as xylene or toluene. The presence of residual salt forms of compounds of formula 1, e.g. as the result of resolution using a chiral resolving agent, do not impede the efficiency of the process.

25 A preferred embodiment of the invention is the racemisation of bupivacaine (1,  $R^1 = n$ -butyl,  $R^2 = 2,6$ -dimethylphenyl) enriched in one enantiomer, preferably the (*R*)-enantiomer, by heating with propanoic acid or butanoic acid. A suitable cosolvent such as xylene allows the  
30 reaction to be conducted at optimum temperature, i.e. about 130°C. Compared to the prior art, this process, and processes of the invention in general, afford significant advantages since neither dilute aqueous solutions ( $[1] < 50$  mM) nor extended reactions times are required.

35 As a further feature of the invention, it has been discovered that asymmetric transformation of the N-heterocyclic-2-carboxanilides can be achieved by including

a chiral acid resolving agent as an additional component in the processes described above (exemplified in Scheme 2). Two variants of such transformations are possible: firstly, a one-pot process in which a given enantiomer of the carboxanilide is converted to its optical antipode by heating to effect racemisation, followed by addition of a chiral acid, resulting in diastereoselective crystallisation of a salt; and secondly, the use of pre-formed racemic carboxanilide as a starting material. Any suitable chiral acid can be used; examples include L- and D-tartaric acid, and *O,O*-dibenzoyl and *O,O*-ditoluoyl derivatives thereof; (*R*)- and (*S*)-10-camphorsulphonic acid; (*R*)- and (*S*)-mandelic acid; (*R*)- and (*S*)-malic acid; (*R*)- and (*S*)-1,1'-binaphthyl-2,2'-diyl hydrogen phosphate; and abietic acid.


An important aspect of this invention relates to the ability to operate the process on an industrial scale. This in turn means that the optically-enriched carboxanilides themselves, e.g. obtained by resolution but to an extent that the predominant enantiomer is insufficiently enantiopure for immediate use, become useful products. This applies to mixtures of enantiomers in which one, usually the (*R*)-enantiomer, is present in an enantiomeric excess of 20 to 80%, preferably 25 to 75%, more preferably 30 to 70%, and most preferably 35 to 65%, with respect to its optical antipode. For example, a mixture enriched in (*R*)-bupivacaine can be used practically, by racemisation of the mixture and subsequent resolution.

The following Example illustrates the invention.


#### Example

A stirred mixture of (*S*)-bupivacaine (0.140 g, 0.49 mmol) and propanoic acid (3.5 ml) was heated to reflux under a nitrogen atmosphere for 7 hours. The resulting solution was cooled and then poured into a mixture of distilled water (20 ml) and ethyl acetate (20 ml). Aqueous ammonia (28% w/v) was added until the pH of the aqueous


layer was 10. The organic layer was separated and the aqueous layer extracted with ethyl acetate (20 ml). The combined organic extracts were washed with distilled water (20 ml), dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure to give racemic bupivacaine (0.137 g, 98%).



Scheme 1




Scheme 2



CLAIMS

1. A process for the racemisation of an optically-enriched chiral N-containing heterocyclic compound having 3 to 7 ring atoms, and a 2-carboxanilide group, which  
5 comprises heating the compound in the presence of an acid selected from alkanolic acids, arylalkanoic acids and acids having a pKa of -1 to +6, relative to water.
2. A process according to claim 1, wherein the compound is a piperidine-2-carboxanilide.
- 10 3. A process according to claim 1, wherein the compound is of formula (1)



- 20 wherein R<sup>1</sup> is H or a substituent of up to 20 C atoms and R<sup>2</sup> is alkyl or aryl of up to 20 C atoms.
4. A process according to claim 3, wherein R<sup>2</sup> is C<sub>6-20</sub> aryl.
5. A process according to claim 4, wherein R<sup>1</sup> is C<sub>1-6</sub> alkyl and R<sup>2</sup> is phenyl optionally substituted with one or  
25 more C<sub>1-4</sub> alkyl groups.
6. A process according to claim 5, wherein the starting material is optically-enriched bupivacaine, i.e. R<sup>1</sup> is n-butyl and R<sup>2</sup> is 2,6-dimethylphenyl.
7. A process according to claim 6, wherein the starting  
30 material is enriched in the (R)-enantiomer.
8. A process according to claim 5, wherein R<sup>1</sup> is H, methyl, n-propyl or cyclopropyl and R<sup>2</sup> is 2,6-dimethylphenyl.
9. A process according to any preceding claim, wherein  
35 the racemisation is carried out in a solution of the acid, neat or mixed with an inert cosolvent.



10. A process according to claim 9, wherein a solution of optically-enriched bupivacaine is heated in propanoic acid or butanoic acid.
- 5 11. A process for the asymmetric transformation of a compound as defined in any of claims 1 to 8, which comprises heating the compound in the presence of an acid as defined in claim 1, a chiral acid resolving agent and an inert cosolvent.
- 10 12. A process according to claim 11, wherein the compound, optically-enriched in a given enantiomer, is converted to its optical antipode.
13. A process according to claim 11, wherein racemic compound is transformed.
- 15 14. A process according to any preceding claim, wherein the acid is an alkanoic or arylalkanoic acid.
15. A process according to any of claims 1 to 13, wherein the acid has the given pKa.
- 20 16. A mixture of enantiomers of a compound as defined in any preceding claim, in which one enantiomer is in an excess of 20 to 80%, e.g. 30 to 70%, with respect to the other enantiomer.
17. A mixture according to claim 16, wherein said one enantiomer is the (R)-enantiomer.
- 25 18. Use of a mixture according to claim 16 or claim 17, for the manufacture of levobupivacaine, by racemisation of the mixture and subsequent resolution.

# INTERNATIONAL SEARCH REPORT

Interr. al Application No  
PCT/GB 95/02247

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07D211/60 C07B55/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07D C07B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 239 710 (ASTRA LÄKEMEDEL AKTIEBOLAG) 7 October 1987 see claims; example 1 ---	1-18
A	JOURNAL OF MEDICINAL CHEMISTRY, vol. 14, no. 9, September 1971 WASHINGTON US, pages 891-892, BENJAMIN F. TULLAR 'Optical isomers of mepivacaine and bupivacaine' see the whole document -----	1-18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

15 November 1995

Date of mailing of the international search report

22. 11. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Henry, J

# INTERNATIONAL SEARCH REPORT

Intern. Application No  
PCT/GB 95/02247

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07D211/60 C07B55/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07D C07B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 239 710 (ASTRA LÄKEMEDEL AKTIEBOLAG) 7 October 1987 see claims; example 1 ---	1-18
A	JOURNAL OF MEDICINAL CHEMISTRY, vol. 14, no. 9, September 1971 WASHINGTON US, pages 891-892, BENJAMIN F. TULLAR 'Optical isomers of mepivacaine and bupivacaine' see the whole document -----	1-18

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

15 November 1995

Date of mailing of the international search report

22. 11. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Henry, J

# INTERNATIONAL SEARCH REPORT

Intern al Application No  
PCT/GB 95/02247

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0239710	07-10-87	SE-B- 451840	02-11-87
		AU-B- 592392	11-01-90
		AU-B- 6644986	09-07-87
		CA-A- 1276154	13-11-90
		IE-B- 59061	15-12-93
		JP-B- 7000598	11-01-95
		JP-A- 62158251	14-07-87
		KR-B- 9408302	12-09-94
		SE-A- 8600017	04-07-87
		US-A- 4870086	26-09-89
-----			